Recombinant Human OFQ/N protein (His Tag)

Catalog Number: PDEH100805

Note: Centrifuge before opening to ensure complete recovery of vial contents.

SpeciesHumanSourceE.coli-derived Human PNOC protein Ser20-Val176, with an N-terCalculated MW17.2 kDaObserved MW23 kDaAccessionQ13519Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	
Calculated MW17.2 kDaObserved MW23 kDaAccessionQ13519Bio-activityNot validated for activityPropertiesPurity>95% as determined by reducing SDS-PAGE.	
Observed MW23 kDaAccessionQ13519Bio-activityNot validated for activityPropertiesPurity>95% as determined by reducing SDS-PAGE.	minal His
AccessionQ13519Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.	
Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.	
Properties Purity > 95% as determined by reducing SDS-PAGE.	
Purity > 95% as determined by reducing SDS-PAGE.	
Endotoxin < 10 EU/mg of the protein as determined by the LAL method	
Storage Generally, lyophilized proteins are stable for up to 12 months wh	nen stored at -20 to -80
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7	days. Aliquots of
reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping This product is provided as lyophilized powder which is shippe	d with ice packs.
Formulation Lyophilized from a 0.2 µm filtered solution in PBS with 5% Treb	nalose and 5%
Mannitol.	
Reconstitution It is recommended that sterile water be added to the vial to prep	are a stock solution of
0.5 mg/mL. Concentration is measured by UV-Vis.	



KDa	М		R	
80 60	11			
40				
30				
20				
12				
· ,	1	1		

> 95 % as determined by reducing SDS-PAGE.

Background

This gene encodes a preproprotein that is proteolytically processed to generate multiple protein products. These products include nociceptin, nocistatin, and orphanin FQ2 (OFQ2). Nociceptin, also known as orphanin FQ, is a 17-amino acid neuropeptide that binds to the nociceptin receptor to induce increased pain sensitivity, and may additionally regulate body temperature, learning and memory, and hunger. Another product of the encoded preproprotein, nocistatin, may inhibit the effects of nociceptin.