

Recombinant Human IL1R2/CD121b Protein (His Tag)

Catalog Number: PKSH031865

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human IL1R2/CD121b protein Met 1-Glu 343, with an C-terminal His
Calculated MW	39.3 kDa
Observed MW	50-55 kDa
Accession	NP_004624.1
Bio-activity	1. Measured by its ability to inhibit IL1 β -dependent proliferation in D10.G4. 1 mouse helper T cells. The ED ₅₀ for this effect is typically 5. 5 μ g/mL in the presence of 40 pg/mL of rhIL1 β . 2. Measured by its ability to inhibit Interferon gamma secretion by human natural killer lymphoma NK-92 cells in the presence of 500pg/mL IL1b. The EC ₅₀ for this effect is typically 20-80ng/mL.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Measured by its ability to inhibit Interferon gamma secretion by human natural killer lymphoma NK-92 cells in the presence of 500pg/mL IL1b. The EC₅₀ for this effect is typically 20-80ng/mL.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Interleukin 1 receptor; type II (IL1R2) also known as CD121b (Cluster of Differentiation 121b) is a cytokine receptor that belongs to the interleukin-1 receptor family. This protein binds interleukin alpha (IL1A); interleukin beta (IL1B); and interleukin 1 receptor; type I (IL1R1/IL1RA); and acts as a decoy receptor that inhibits the activity of its ligands. The pleiotropic cytokine IL1 is produced to regulate development and maintenance of the inflammatory responses; and binds to specific plasma membrane receptors on cells. Two distinct types of IL1 receptors which are able to bind IL1 specifically have been identified; designated as IL1RI (IL1RA) and IL1RII (IL1RB). IL1R1 contributes to IL-1 signaling; whereas the IL-1R2/CD121b has no signaling property and acts as a decoy for IL-1. IL-1R2/CD121b structurally consisting of a ligand binding portion comprised of three Ig-like domains; a single transmembrane region; and a short cytoplasmic domain; is expressed in a variety of cell types including B lymphocytes; neutrophils; monocytes; large granular leukocytes and endothelial cells. Interleukin 4 (IL4) is reported to antagonize the activity of interleukin 1 by inducing the expression and release of this cytokine.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017