# Recombinant Human CD3E&CD3G Heterodimer (C-Fc-6His&C-Fc-Flag)

Catalog Number: PKSH034015



Note: Centrifuge before opening to ensure complete recovery of vial contents.

				. •	
	00	cri	m	-10	m
JU	$\mathbf{c}$		174	7 U	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
			т.		

 Species
 Human

 Mol\_Mass
 39.2&38.1 kDa

 Accession
 P07766&P09693

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

#### Background

T-cell surface glycoprotein CD3 epsilon & CD3 gamma chain, also known as CD3E & CD3G, are single-pass type I membrane proteins. CD3D, together with CD3- epsilon(CD3E), CD3-gamma and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex T cell receptor-CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways.

## For Research Use Only