

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® Red 780 Anti-Mouse CD16/32 Antibody[2.4G2]

Catalog Number: E-AB-F0997S

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Rat Host

Isotype Rat IgG2b, ĸ Clone No. 2.4G2

Isotype Control Elab Fluor® Red 780 Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09842S]

Conjugation Elab Fluor®Red 780

Conjugation Information Elab Fluor® Red 780 is designed to be excited by the Red (627-640 nm) laser and

detected using an optical filter centered near 770 nm (e.g., a 780/60 nm bandpass filter).


Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



C57BL/6 murine splenocytes are stained with Elab Fluor® Red 780 Anti-Mouse CD16/32 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD16a/b;CD32;CD32A/B;FCG2A;FCGR2A/BFCGR3;FCGR3A/B;Fc fragment of IqG low

affinity Illa/b receptor;Fc fragment of IgG low affinity Illb receptor;Fc fragment of IgG low

affinity IIa/b receptor;Fc gamma RIIa/bFc gamma receptor III A/B;FcGR

Uniprot ID P08508;P08101

For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com Web:www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID

Background

14130;14131

CD16 is low affinity IgG Fc receptor III (FcR III) and CD32 is FcR II. CD16/CD32 are expressed on B cells, monocytes/macrophages, NK cells, granulocytes, mast cells, and dendritic cells. The Fc receptors bind antibody-antigen immune complexes and mediate adaptive immune responses.