

## Purified Anti-Mouse CD48 Antibody[HM48-1]

catalog number: E-AB-F1017A

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Mouse
<b>Host</b>	Armenian Hamster
<b>Isotype</b>	Armenian Hamster IgG
<b>Clone</b>	HM48-1
<b>Buffer</b>	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

### Applications Recommended Dilution

<b>FCM</b>	2 µg/mL( $1 \times 10^5$ - $5 \times 10^5$ cells)
------------	---

### Data



C57/BL6 Mouse splenocytes were stained with 0.2 µg Purified Anti-Mouse CD48 Antibody[HM48-1] (Right) and 0.2 µg Armenian Hamster IgG, κ Isotype Control (Left), followed by PE-conjugated Goat Anti-Armenian Hamster IgG Secondary Antibody.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

CD48 is a 45 kD GPI-anchored glycoprotein also known as BCM1, Blast-1 (human), and OX-45 (rat). It is a member of the Ig superfamily, expressed on T and B cells and monocytes/macrophages. It plays a role in adhesion and T cell recognition. The primary ligands for CD48 are CD2 and CD244.

### For Research Use Only