

PE Anti-Mouse LPAM-1 Antibody[DATK32]

Catalog Number: GFH00563D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Mouse
Host	Rat
Isotype	Rat IgG2a, κ
Clone No.	DATK32
Isotype Control	PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832D]
Conjugation	PE
Conjugation Information	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide.


Applications

FCM

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes cells with APC Anti-Mouse B220 Antibody and PE Anti-Mouse LPAM-1 Antibody[DATK32] (left) or PE Rat IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	$\alpha 4\beta 7$ Integrin;CD49d/ $\beta 7$;LPAM-1;ITGA4;ITGB7
Uniprot ID	P26011;Q00651
Gene ID	311144;25713

For Research Use Only

Background

DATK32 antibody is specific for a combinatorial determinate of integrin $\alpha 4\beta 7$ complex. Integrin $\alpha 4\beta 7$ is composed of a 150 kD ($\alpha 4$ or CD49d) and a 130 kD ($\beta 7$) heterodimer, also known as CD49d/ $\beta 7$ or LPAM-1. Belonging to the Ig superfamily, it is found on the majority of peripheral lymphocytes and subsets of thymocytes and bone marrow cells (including mast cell progenitors). Integrin $\alpha 4\beta 7$ binds its ligands, VCAM-1 (CD106), MAdCAM-1 and fibronectin, and plays an important role in lymphocytes adhesion and the direction of migration of blood lymphocytes to the intestine and associated lymphoid tissues.