A Reliable Research Partner in Life Science and Medicine

CK-18 Polyclonal Antibody

catalog number: E-AB-70095

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen KLH conjugated Synthetic peptide corresponding to Mouse Cytokeratin 18

Host Rabbit
Isotype IgG

Purification Affinity purification


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein

protectant and 50% glycerol.

Applications	Recommended	Dilution
Tippiications	1 C COMMITTE HUCU	Dilution


WB 1:500-1:2000 IHC 1:200-1:800

Data




Western Blot analysis of various samples using CK-18 Polyclonal Antibody at dilution of 1:1000.


Observed-MW:48 kDa Calculated-MW:48 kDa




Immunohistochemistry analysis of paraffin-embedded human stomach using CK-18 Polyclonal Antibody at dilution of



Immunohistochemistry analysis of paraffin-embedded Rat lung using CK-18 Polyclonal Antibody at dilution of 1:500.



Immunohistochemistry analysis of paraffin-embedded human kidney using CK-18 Polyclonal Antibody at dilution of 1:500.



Immunohistochemistry analysis of paraffin-embedded Human Placenta using CK-18 Polyclonal Antibody(Elabscience Product Detected by Lifespan).

For Research Use Only

Fax: 1-832-243-6017

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

Cytokeratin 18 (CK18) has a molecular weight of 45 KDa, which is a low molecular weight type A cytokeratin. It is a marker of simple epithelium, including glandular epithelium. The stratified squamous epithelium is generally negative. CK18 is mainly used in the diagnosis of adenocarcinoma.

Fax: 1-832-243-6017