

Purified Anti-Human CD45 Antibody[HI30]

catalog number: E-AB-F1137A

Note: Centrifuge before opening to ensure complete recovery of vial contents.



Description

Reactivity	Human
Immunogen	Recombinant Human CD45 protein
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	HI30
Purification	>98%, Protein A/G purified
Conjugation	Unconjugated
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

FCM	2 $\mu\text{g/mL}$ (0.5×10^6 - 1×10^6 cells)
IHC	1:50-1:100

Data



Human peripheral blood were stained with 0.2 μg Purified Anti-Human CD45 Antibody[HI30] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Immunohistochemistry of paraffin-embedded Human tonsil using CD45 Monoclonal Antibody at dilution of 1:100.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

CD45 is a 180 - 240 kD single chain type I membrane glycoprotein also known as leukocyte common antigen (LCA) and T200. It is a tyrosine phosphatase expressed on the plasma membrane of all hematopoietic cells, except erythrocytes or platelets. CD45 is a signaling molecule that regulates a variety of cellular processes including cell growth, differentiation, cell cycle, and oncogenic transformation. CD45 plays a critical role in T and B cell antigen receptor-mediated activation by dephosphorylating substrates including p56Lck, p59Fyn, and other Src family kinase s. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to bind galectin-1 and to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4.