Recombinant Human GDF5/BMP-14 Protein

Catalog Number: PKSH033660

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	E.coli-derived Human GDF5; BMP-14 protein Ala382-Arg501, with an C-terminal His
Calculated MW	14.5 kDa
Observed MW	18 kDa
Accession	P43026
Bio-activity	Measure by its ability to induce alkaline phosphatase production by ATDC5 cells.The
	ED_{50} for this effect is <14 ng/mL.
Properties	
Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $<$ -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20 mM sodium citrate, 0.2 M NaCl, pH 3.5.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Web:www.elabscience.com

Elabscience®

Growth Differentiation Factor 5(GDF-5, BMP-14) is a member of the BMP family of TGFβ superfamily proteins. Human GDF-5, -6, and -7 are a defined subgroup of the BMP family. GDF-5 is synthesized as a homodimeric precursor protein consisting of a 354 amino acid (aa) Nterminal proregion and a 120 aa C-terminal mature peptide. Mature human GDF-5 shares 99% aa sequence identity with both mature mouse and rat GDF-5. GDF-5 signaling is mediated by formation of a heterodimeric complex consisting of a type 1 (BMPR-IB) and a type II (BMPR-IIor Activin RII) serine/threonine kinase receptor which results in the phosphorylation and activation of cytosolic Smad proteins (Smad1, 5, and 8). GDF-5 is involved in multiple developmental processes including limb generation, cartilage development, joint formation, bone morphogenesis, cell survival, and neuritogenesis. Inhibition of GDF-5 expression or alteration of its signaling can facilitate the development of osteoarthritis.