

## Biotin Anti-Mouse TCR $\gamma/\delta$ Antibody[GL3]

**Catalog Number:** E-AB-F1282B

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

<b>Reactivity</b>	Mouse
<b>Host</b>	Armenian Hamster
<b>Isotype</b>	Armenian Hamster IgG
<b>Clone No.</b>	GL3
<b>Isotype Control</b>	Biotin Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F09853B]
<b>Conjugation</b>	Biotin
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per $10^6$ cells in $100 \mu\text{L}$ volume or $100 \mu\text{L}$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
------------	---

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at $2-8^{\circ}\text{C}$ for 12 months. Do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	T cell receptor $\gamma/\delta$
<b>Uniprot ID</b>	Q96E93;O88713
<b>Gene ID</b>	110066;110067
<b>Background</b>	T cell receptor (TCR) is a heterodimer consisting of an $\alpha$ and a $\beta$ chain (TCR $\alpha/\beta$ ) or a $\gamma$ and a $\delta$ chain (TCR $\gamma/\delta$ ). TCR $\gamma/\delta$ belongs to the immunoglobulin superfamily, which is involved in the recognition of certain bacterial and tumor antigens bound to MHC class I. $\gamma/\delta$ TCR associates with CD3 and is expressed on a T cell subset found in the thymus, the intestinal epithelium, and the peripheral lymphoid tissues and peritoneum. Most $\gamma/\delta$ T cells are CD4-/CD8- although some are CD8+. T cells expressing the $\gamma/\delta$ TCR have been shown to play a role in oral tolerance, tumor-associated tolerance, and autoimmune disease. It has been reported that $\gamma/\delta$ T cells also play a principal role in antigen presentation.

### For Research Use Only