

Purified Anti-Mouse CD71 Antibody[R17 217.1.3/TIB-219], Functional Grade

catalog number: E-AB-F10930

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description



Reactivity	Mouse
Immunogen	Recombinant Mouse CD71 protein
Host	Rat
Isotype	Rat IgG2a, κ
Clone	R17 217.1.3/TIB-219
Purification	>98%, Protein A/G purified
Buffer	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

Applications

Recommended Dilution

FCM	$\leq 0.2 \mu\text{g}$ per million cells in 100 μL volume
Depletion	Reported in the literature

Data



C57/BL6 Mouse bone marrow cells were stained with 0.2 μg Purified Anti-Mouse CD71 Antibody[R17 217.1.3/TIB-219], Functional Grade (Right) and 0.2 μg Rat IgG2a, κ

Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody, then anti-Mouse Ter-119 PE-conjugated Monoclonal Antibody.

C57/BL6 Mouse bone marrow cells were stained with 0.2 μg Purified Anti-Mouse CD71 Antibody[R17 217.1.3/TIB-219], Functional Grade (Right) and 0.2 μg Rat IgG2a, κ

Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody, then anti-Mouse Ter-119 PE-conjugated Monoclonal Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V3.0

CD71 is a 95 kD type II heterodimeric transmembrane glycoprotein that is also known as T9 and transferrin receptor. CD71 is expressed on proliferating cells, reticulocytes, and erythroid precursors. Its expression is very low on resting leukocytes. CD71 plays a role in the control of cellular proliferation by facilitating the uptake of iron via ferrotransferrin binding and the recycling of apotransferrin to the cell surface.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

Application References

Natalia Torow, et al. Nat Commun. 2015 Jul 21;6:7725.