CYTIP Polyclonal Antibody

catalog number: E-AB-53171



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Fusion protein of human CYTIP
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
IHC	1:100-1:300
Data	
cancer tissue using CYT	of paraffin-embedded Human gastric Immunohistochemistry of paraffin-embedded Human IP Polyclonal Antibody at dilution of esophagus cancer tissue using CYTIP Polyclonal Antibody 1:100(×200) at dilution of 1:100(×200)
Preparation & Storage	1.100(~200) at dilution of 1.100(~200)
	Store at 2000 Malid for 12 months. Assaid frages / them evalue
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.
Background	
The metric area ded her this zero contains 2 leaving size or domains and a putation C terminal method to size of	

The protein encoded by this gene contains 2 leucine zipper domains and a putative C-terminal nuclear targeting signal, but does not have any hydrophobic regions. This protein is expressed weakly in resting NK and T cells. The encoded protein modulates the activation of ARF genes by CYTH1. This protein interacts with CYTH1 and SNX27 proteins and may act to sequester CYTH1 protein in the cytoplasm.

For Research Use Only