Elabscience®

PTPA Polyclonal Antibody

catalog number: E-AB-53148

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human;Mouse	
Immunogen	Fusion protein of human PTPA	
Host	Rabbit	
Isotype	IgG	
Purification	Antigen affinity purification	
Conjugation	Unconjugated	
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilution	
IHC	1:50-1:200	
Data		
Immunohistochemistry of	f paraffin-embedded Human ovarianImmunohistochemistry of paraffin-embedded Human	
cancer tissue using PTPA Polyclonal Antibody at dilution of thyroid cancer tissue using PTPA Polyclonal Antibody at		

cancer tissue using PTPA Polyclonal Antibody at dilution of thyroid cancer tissue using PTPA Polyclonal Antibody at 1:60(×200)

dilution of 1:60(×200)

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the
	temperature recommended.

Background

Protein phosphatase 2A is one of the four major Ser/Thr phosphatases and is implicated in the negative control of cell growth and division. Protein phosphatase 2A holoenzymes are heterotrimeric proteins composed of a structural subunit A, a catalytic subunit C, and a regulatory subunit B. The regulatory subunit is encoded by a diverse set of genes that have been grouped into the B/PR55, B'/PR61, and B"/PR72 families. These different regulatory subunits confer distinct enzymatic specificities and intracellular localizations to the holozenzyme. The product of this gene belongs to the B' family. This gene encodes a specific phosphotyrosyl phosphatase activator of the dimeric form of protein phosphatase 2A. Alternative splicing results in multiple transcript variants encoding different isoforms.