

# Purified Anti-Mouse CD25 Antibody[2E4]

catalog number: AN006190P

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Mouse
<b>Immunogen</b>	Recombinant Mouse CD25 protein
<b>Host</b>	Rabbit
<b>Isotype</b>	Rabbit IgG, $\kappa$
<b>Clone</b>	2E4
<b>Purification</b>	>98%, Protein A/G purified
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	PBS, pH 7.2. Contains 0.05% proclin 300.

## Applications

## Recommended Dilution

<b>FCM</b>	2 $\mu\text{g}/\text{mL}$ ( $1 \times 10^5$ - $5 \times 10^5$ cells)
------------	--

## Data



C57/BL6 Mouse splenocytes were stained with 0.2  $\mu\text{g}$  Purified Anti-Mouse CD25 Antibody[2E4] (Right) and 0.2  $\mu\text{g}$  Rabbit IgG,  $\kappa$  Isotype Control (Left), followed by Alexa Fluor<sup>®</sup> 647-conjugated Goat Anti-Rabbit IgG Secondary Antibody, then anti-Mouse CD4 FITC-conjugated Monoclonal Antibody.

## Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	Order now, ship in 3 days

## Background

Receptor for interleukin-2. The receptor is involved in the regulation of immune tolerance by controlling regulatory T cells (TREGs) activity. TREGs suppress the activation and expansion of autoreactive T-cells.

## For Research Use Only