



A Reliable Research Partner in Life Science and Medicine

Rat FAS/CD95 Antibody Pair Set

Catalog No. E-KAB-0379 Applications ELISA

Synonyms FASR, TNFRSF6, APO1, APT1, FAS1, FASTM, ALPS1A

Kit components & Storage

Title	Specifications	Storage
Rat FAS/CD95 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Rat FAS/CD95 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0379)	
		Rat FAS/CD95 Capture Antibody	Rat FAS/CD95 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Rat FAS/CD95 protein	Recombinant Rat FAS/CD95 protein
Information	Swissprot	Q63199	
Product details	Reactivity	Rat	Rat
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Rat FAS/CD95 in ELISAs.	·

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

Applications

Rat FAS/CD95 Sandwich ELISA Assav:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Rat FAS/CD95 Capture Antibody	
Capture			Ais 1
ELISA Detection	1:1000-1:10000	Rat FAS/CD95 Detection Antibody (Biotin)	Optical Density
			0.01 100 1000 10000 Rat FAS/CD95 concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

FAS,also named as CD95,APO-1,APT1,FAS1 and TNFRSF6,is a receptor for TNFSF6/FASLG. It is a cell surface receptor belonging to the TNF receptor superfamily,can mediates apoptosis by ligation with an agonistic anti-Fas antibody or Fas ligand. Stimulation of Fas results in the aggregation of its intracellular death domains,leading to the formation of the death-inducing signaling complex (DISC). FAS-mediated apoptosis may have a role in the induction of peripheral tolerance,in the antigen-stimulated suicide of mature T-cells,or both. The secreted isoforms 2 to 6 block apoptosis (in vitro). This anti-Fas monoclonal antibody can be used to induce apoptosis in cell cultures through Fas by imitating the Fas-ligand.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com