

Elab Fluor® Violet 450 Anti-Human IL-12/IL-23 p40 Antibody[C8.6]

Catalog Number: AN00843Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	C8.6
Isotype Control	Elab Fluor® Violet 450 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792Q]
Conjugation	Elab Fluor® Violet 450
Conjugation Information	Elab Fluor® Violet 450 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

Data



IFN- γ -primed and then LPS-stimulated (24h) human peripheral blood monocytes intracellular stained with Elab Fluor® Violet 450 Anti-Human IL-12/IL-23 p40 Antibody[C8.6] (filled gray histogram) or Elab Fluor® Violet 450 Mouse IgG1, κ Isotype Control (empty black histogram).

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protect from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	Interleukin-12 p40;Interleukin-23 p40;Cytotoxic lymphocyte maturation factor (CLMF);Natural killer cell stimulatory factor (NKSF);CTL maturation factor (TcMF);T-cell stimulating factor (TSF)
------------------------	--

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3

Uniprot ID P29460

Gene ID 3593

Background
The C8.6 antibody reacts with human IL-12 p40 subunit of the IL-12 p70 heterodimer and IL-23 p40 subunit of the IL-23 p19/p40, as well as p40 monomer and homodimer. The C8.6 antibody has been reported to strongly inhibit different biological activities of IL-12, (e.g., IFN-gamma induction, mitogenic effects on PHA blasts, and enhancement of NK cell-mediated cytotoxicity). The C8.6 antibody can neutralize the bioactivity of natural or recombinant IL-12.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3