

Recombinant Mouse NAALADL1 Protein (His Tag)

Catalog Number: PKSM040428

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|--|
| Species | Mouse |
| Source | HEK293 Cells-derived Mouse NAALADL1 protein Pro 29-Leu 745, with an N-terminal His |
| Calculated MW | 80.0 kDa |
| Observed MW | 110 kDa |
| Accession | Q7M758 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|---|
| Purity | > 97 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 97 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

N-acetylated-alpha-linked acidic dipeptidase-like protein, also known as NAALADL1, NAALADase L, and Ileal dipeptidylpeptidase, is a Single-pass type II membrane protein and a member of the peptidase M28 family and M28B subfamily. NAALADase L is mainly expressed in the distal small intestine. It is also expressed in the spleen and testis and Weakly expressed in the brain, locating mainly to the brain stem, amygdala, thalamus and ventral striatum. NAALADase L is a chloride-activated, membrane bound, metallopeptidase that cleaves the endogenous neuropeptide N-acetyl-aspartyl-glutamate (NAAG). NAAG acts as a partial NMDA agonist as well as a full agonist at the presynaptic metabotropic glutamate receptor 3 (mGluR3), where it acts to reduce glutamate release. NAALADase L also exhibits a dipeptidyl-peptidase IV type activity. NAALADase inhibition may be a novel therapeutic approach to reduce or inhibit heightened aggressiveness, and possibly to treat aggressive behavior associated with psychiatric disorders.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017