

## Recombinant Human GSTAL Protein(Trx Tag)

**Catalog Number:** PDEH100656

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human GSTAL protein Met1-Phe222, with an N-terminal Trx
<b>Calculated MW</b>	44.4 kDa
<b>Observed MW</b>	39 kDa
<b>Accession</b>	P08263
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Human GSTAL proteins, 2µg/lane of  
Recombinant Human GSTAL proteins was resolved with  
SDS-PAGE under reducing conditions, showing bands at 39  
KD

### Background

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Rev. V1.2

GSTA1 (Glutathione S-Transferase Alpha 1) is a Protein Coding gene. This gene encodes a member of a family of enzymes that function to add glutathione to target electrophilic compounds. Glutathione S-transferases (GSTs) are involved in the detoxification of carcinogens and may be linked to carcinogenesis. As a vital component of GSTs, GSTA1 plays an important role in carcinogenesis. GSTA1 expression may be a target molecule in the early diagnosis and treatment of lung cancer. Human colonic adenocarcinoma (Caco-2) cells in culture undergo spontaneous differentiation into mature enterocytes in association with progressive increases in expression of glutathione S-transferase alpha-1 (GSTA1). GSTA1 levels may play a role in modulating enterocyte proliferation but do not influence differentiation or apoptosis. GSTA1 may play a key role during pregnancy.