

A Reliable Research Partner in Life Science and Medicine

# FITC Anti-Human CD11a Antibody[R7-1]

Catalog Number: E-AB-F1211C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human Mouse Host

Mouse IgG1, κ Isotype

Clone No. R7-1

FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792C] Isotype Control

Conjugation

**Conjugation Information** FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

#### **Applications** Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



Human peripheral blood lymphocytes are stained with FITC Anti-Human CD11a Antibody (filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

### **Preparation & Storage**

Keep as concentrated solution. **Storage** 

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

#### **Antigen Information**

**Alternate Names** CD11A;ITGAL;Integrin alpha-L;LFA-1A

**Uniprot ID** P20701 Gene ID 3683

## For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

### **Background**

Elabscience®

CD11a is a 170-180 kD type I transmembrane glycoprotein also known as LFA-1a chain and integrin αL subunit. CD11a non-covalently associates with integrin β2 (CD1 8) to form LFA-1. It is expressed on all leukocytes, including B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. It is absent on nonhematopoietic tissues and platelets. CD11a plays a central role in leukocyte cell-cell interactions and is important in lymphocyte costimulation. CD11a/CD18 binds to ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Email:techsupport@elabscience.com Web:www.elabscience.com