



A Reliable Research Partner in Life Science and Medicine

# Purified Anti-Mouse CD205 Antibody[NLDC-145]

catalog number: AN008440P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Mouse

Immunogen Recombinant Mouse CD205 protein

**Host** Ra

**Isotype** Rat WI, also known as Wistar (outbred) IgG2a, κ

Clone NLDC-145

**Purification** >98%, Protein A/G purified


**Buffer** Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

## Applications Recommended Dilution

**FCM** 2  $\mu$ g/mL(0.5×10<sup>6</sup>-1×10<sup>6</sup> cells)

#### Data



A20 were stained with 0.2 μg Purified Anti-Mouse CD205 Antibody[NLDC-145] (Right) and 0.2 μg Rat IgG2a, κ Isotype Control(Left), followed by APC-conjugated Goat Anti-Rat IgG Secondary Antibody.

## **Preparation & Storage**

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

### **Background**

CD205, also known as DEC-205, is a 205 kD integral membrane protein homologous to the macrophage mannose receptor. It is a type I cell surface protein that belong to the C-type lectin family. CD205 is expressed at high levels by dendritic cells and thymic epithelial cells. It is also expressed by a number of other cell types, such as B lymphocytes, macrophages, Langerhans cells, bone marrow stromal cells, granulocytes, epithelial cells of pulmonary airways, and the capillaries of the brain. CD205 is a novel endocytic receptor used by dendritic cells and thymic epithelial cells to direct captured antigens from the extracellular space to specialized antigen processing. It mediates antigen uptake and presentation and cross-presentation to T cells. It has been reported that CD205 acts as a recognition receptor for dying cells, potentially provides an important pathway for the uptake of self-antigen in the intrathymic environment, and is involved in peripheral tolerance. Antibody-mediated antigen-targeting via the DEC-205 receptor increases the efficiency of vaccination for T cell immunity.

For Research Use Only