A Reliable Research Partner in Life Science and Medicine

# Recombinant Phospho-Stat5 (Tyr694) Monoclonal Antibody

catalog number: AN300135L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Human

**Immunogen** A synthetic phosphopeptide corresponding to residues around Tyr694 of human

Phospho-Stat5.


Host Rabbit
Isotype IgG
Clone 3G5
Purification Protein A

Buffer 10 mM sodium HEPES, 150 mM NaCl, 100 μg/mL protein protectant, 50% glycerol, pH

7.5

## **Applications** Recommended Dilution

**WB** 1:1000-1:10000 **IHC-P** 1:50-1:200



Immunohistochemistry of paraffin-embedded human breast cancer using Phospho-Stat5 (Tyr694) Monoclonal Antibody at dilution of 1:50.

## Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

### Background

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L2/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. Alternatively spliced transcript variants have been found for this gene.

# For Research Use Only