

Recombinant Phospho-Stat5 (Tyr694) Monoclonal Antibody

catalog number: **AN300135L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	A synthetic phosphopeptide corresponding to residues around Tyr694 of human Phospho-Stat5.
Host	Rabbit
Isotype	IgG
Clone	3G5
Purification	Protein A
Buffer	10 mM sodium HEPES, 150 mM NaCl, 100 µg/mL protein protectant, 50% glycerol, pH 7.5


Applications

Applications	Recommended Dilution
WB	1:1000-1:10000
IHC-P	1:50-1:200

Data




Immunohistochemistry of paraffin-embedded human lung cancer using Phospho-Stat5 (Tyr694) Monoclonal Antibody at dilution of 1:50.




Western blot analysis of extracts from serum-starved TF-1, untreated (-) or treated with GM-CSF (Cat: PKSH031982) (100 ng/mL, 5 min; +), using Phospho-Stat5 (Tyr694) Antibody at 1:1000 dilution (upper) or Anti-STAT5a Antibody, Rabbit PAb, Antigen Affinity Purified at 1:500 dilution (lower).

Observed-MW: 100 kDa
Calculated-MW: 90 kDa




Immunohistochemistry of paraffin-embedded human breast cancer using Phospho-Stat5 (Tyr694) Monoclonal Antibody at dilution of 1:50.



Western blot analysis of extracts from serum-starved TF-1, untreated (line A); treated with GM-CSF (Cat: PKSH031982) (100 ng/mL, 5 min) (line B); treated with GM-CSF and λ-phosphatase (line C) using Phospho-Stat5 (Tyr694) Antibody at 1:1000 dilution.

Observed-MW: 100 kDa
Calculated-MW: 90 kDa

For Research Use Only



Western blot analysis of extracts from serum-starved TF-1, untreated (line A); treated with GM-CSF (Cat: PKSH031982) (100 ng/mL, 5 min; +), without peptide (line B) or antigen-specific phosphopeptide (line C) or antigen-specific peptide (line D) using Phospho-Stat5 (Tyr694) Antibody at 1:1000 dilution.

Observed-MW:100 kDa

Calculated-MW:90 kDa

Preparation & Storage

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping

Ice bag

Background

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L2/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. Alternatively spliced transcript variants have been found for this gene.

For Research Use Only