

FGF-2/bFGF/FGF-b (Gly132-Ser288), Human, Recombinant**Cat. No. : GPCK007****产品信息**

物种	Human
表达宿主	E.coli
序列信息	Gly132-Ser288
检索号	P09038-4
分子量	17.4 kDa
有效期	12 months
生物活性	Measured in a cell proliferation assay using BALB/c 3T3 cells. The ED50 for this effect is 0.42 ng/mL.

产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μm filtered solution of 20 mM Tris, 150 mM NaCl, 3% Trehalose, 4% Mannitol, pH 7.5.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

FGF-basic is a members of the Fibroblast Growth Factors (FGFs) family. The family constitutes a large family of Proteins involved in many aspects of development including cell proliferation, growth, and differentiation. They act on several cell types to regulate diverse physiologic functions including angiogenesis, cell growth, pattern formation, embryonic development, metabolic regulation, cell migration, neurotrophic effects, and tissue repair. FGF-basic is a non-glycosylated heparin binding Growth Factor that is expressed in the brain, pituitary, kidney, retina, bone, testis, adrenal gland liver, monocytes, epithelial cells and endothelial cells. FGF-basic signals through FGFR 1b, 1c, 2c, 3c and 4.

