



A Reliable Research Partner in Life Science and Medicine

Recombinant KLK8/Kallikrein 8 Monoclonal Antibody

catalog number: AN300243P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human KLK8 / Kallikrein 8 Protein


HostRabbitIsotypeIgGCloneB161PurificationProtein A

Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution

WB 1:500-1:1000

Data



Western Blot with KLK8 / Kallikrein 8 Monoclonal Antibody at

dilution of 1:500. Lane A: A431 Whole Cell Lysate,

Lysates/proteins at 30 µg per lane.

Observed-MW:35 kDa Calculated-MW:28 kDa

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when

stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

Kallikrein-8, also known as Neuropsin, Serine protease 19, Serine protease TADG-14, Tumor-associated differentially expressed gene 14 protein, and KLK8 is a secreted protein that belongs to the peptidase S1 family and Kallikrein subfamily. It is a serine protease that is capable of degrading some proteins such as casein, fibrinogen, kininogen, fibronectin, and collagen type IV. Kallikrein-8/KLK8 plays a role in the formation and maturation of orphan and small synaptic boutons in the Schaffer-collateral pathway. It regulates Schaffer-collateral long-term potentiation in the hippocampus and is required for memory acquisition and synaptic plasticity. It is involved in skin desquamation and keratinocyte proliferation and plays a role in the secondary phase of pathogenesis following spinal cord injury. It also cleaves L1CAM in response to increased neural activity. It induces neurite outgrowth and fasciculation of cultured hippocampal neurons. Kallikrein-8/KLK8 is expressed at high levels in serum, ascites fluid, and tumor cytosol of advanced-stage ovarian cancer patients and may serve as a marker of ovarian cancer. Kallikrein-8/KLK8 may have potential clinical value for disease diagnosis or prognosis and it may also be a useful therapeutic target.

For Research Use Only