

Recombinant Cytokeratin 19 Monoclonal Antibody

catalog number: **AN300920L**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Human;Mouse;Rat
Immunogen	Recombinant Human Cytokeratin 19 protein
Host	Rabbit
Isotype	Rabbit IgG
Clone	C110544
Purification	Protein A
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications

Recommended Dilution

IHC	1:400-1:2000
WB	1:1000-1:5000
IF	1:200-1:1000
ELISA	1:5000-1:20000


Data



Western Blot with Recombinant Cytokeratin 19 Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: K562 cells.

Observed-MW:40 kDa

Calculated-MW:40 kDa



Immunohistochemistry of paraffin-embedded rat lungs using Recombinant Cytokeratin 19 Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

The protein encoded by this gene is a member of the keratin family. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. The type I cytokeratins consist of acidic proteins which are arranged in pairs of heterotypic keratin chains. Unlike its related family members, this smallest known acidic cytokeratin is not paired with a basic cytokeratin in epithelial cells. It is specifically expressed in the periderm, the transiently superficial layer that envelopes the developing epidermis. The type I cytokeratins are clustered in a region of chromosome 17q12-q21.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.4