## Recombinant Human CD46 Protein (His Tag)

Catalog Number: PKSH032220



Description		
Species	Human	
Mol_Mass	33.8 kDa	
Accession	P15529-11	
<b>Bio-activity</b>	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Data		
	kDa	MK
	120 90	and the second s
	90 60	- Cence
	10	- Elsa
	30 20	- science'
		Elabsor
	20	
	14	-
		and the second

> 95 % as determined by reducing SDS-PAGE.

## Background

CD46 is a type I membrane protein containing four Sushi domains. CD46 is expressed by all cells except erythrocytes. CD46 has cofactor activity for inactivation of complement components C3b and C4b by serum factor I, which protects the host cell from damage by complement. It may be involved in the fusion of the spermatozoa with the oocyte during fertilization. CD46 also acts as a costimulatory factor for T-cells which induces the differentiation of CD4+ into Tregulatory 1 cells. T-regulatory 1 cells suppress immune responses by secreting interleukin-10, and therefore are thought to prevent autoimmunity. A number of viral and bacterial pathogens exploit this property and directly induce an immunosuppressive phenotype in T-cells by binding to CD46. CD46 acts as a receptor for the Edmonston strain of measles virus, human herpesvirus-6, and type IV pili of pathogenic Neisseria.

## For Research Use Only