

# Recombinant Human STC2/Stanniocalcin 2 Protein (His Tag)



Catalog Number:PKSH030583

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

|                                    |              |
|------------------------------------|--------------|
| <b>Synonyms</b>                    | STC-2;STCRP  |
| <b>Species</b>                     | Human        |
| <b>Expression Host</b>             | HEK293 Cells |
| <b>Sequence</b>                    | Met 1-Arg302 |
| <b>Accession</b>                   | O76061       |
| <b>Calculated Molecular Weight</b> | 32.1 kDa     |
| <b>Observed molecular weight</b>   | 35-42 kDa    |
| <b>Tag</b>                         | C-His        |

## Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | < 1.0 EU per $\mu$ g of the protein as determined by the LAL method.  |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from sterile PBS, pH 7.4<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual.            |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

STC2 is a secreted; homodimeric glycoprotein which expressed in a wide variety of tissues. STC2 has an anti-hypocalcemic action on calcium and phosphate homeostasis. It may have autocrine or paracrine functions. Its C-terminus contains a cluster of histidine residues which may interact with metal ions. STC2 has 10 of its 15 cysteine residues conserved among stanniocalcin family members and is phosphorylated by casein kinase 2 exclusively on its serine residues. It may play a role in the regulation of renal and intestinal calcium and phosphate transport; cell metabolism; or cellular calcium/phosphate homeostasis.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017