Recombinant Human RPLP1 Protein(Trx Tag)

Catalog Number: PDEH100648



Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}					٠,		
H)	es	C1	ľ٦	n	П	on	

Species Human

Source E.coli-derived Human Human RPLP1 protein Met1-Asp114, with an N-terminal Trx

 Mol_Mass
 32.5 kDa

 Accession
 P05386

Bio-activity Not validated for activity

Properties

Purity > 95% as determined by reducing SDS-PAGE.

Endotoxin < 10 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


Shipping This product is provided as lyophilized powder which is shipped with ice packs.
 Formulation Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis

Data



SDS-PAGE analysis of Human RPLP1 proteins, 2µg/lane of Recombinant Human RPLP1 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 35

KD

Background

For Research Use Only

Recombinant Human RPLP1 Protein(Trx Tag)

Catalog Number: PDEH100648



Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal phosphoprotein that is a component of the 60S subunit. The protein, which is a functional equivalent of the E. coli L7/L12 ribosomal protein, belongs to the L12P family of ribosomal proteins. It plays an important role in the elongation step of protein synthesis. Unlike most ribosomal proteins, which are basic, the encoded protein is acidic. Its C-terminal end is nearly identical to the C-terminal ends of the ribosomal phosphoproteins P0 and P2. The P1 protein can interact with P0 and P2 to form a pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein is located in the cytoplasm. Two alternatively spliced transcript variants that encode different proteins have been observed. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.