Elabscience®

(KO Validated) SUCLG2 Polyclonal Antibody

catalog number: E-AB-65605

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human SUCLG2 (NP_003839.2).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:200
IF	1:50-1:100

Data





Western blot analysis of extracts of various cell lines using SUCLG2 Polyclonal Antibody at dilution of 1:3000.

Observed-MW:47 kDa Calculated-MW:46 kDa/47 kDa



Immunohistochemistry of paraffin-embedded Human liver cancer using SUCLG2 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded Human lung cancer using SUCLG2 Polyclonal Antibody at dilution of

1:100 (40x lens).





For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w .elabscience.com</u>

Elabscience®

Immunofluorescence analysis of U-2 OS cells using SUCLG2 Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

This gene encodes a GTP-specific beta subunit of succinyl-CoA synthetase. Succinyl-CoA synthetase catalyzes the reversible reaction involving the formation of succinyl-CoA and succinate. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 5 and 12.

For Research Use Only