

FGF-20/FGFK, Human, Recombinant

Cat. No. : PCK294

General Information

Synonyms	FGFK
Species	Human
Expression host	E.coli
Sequence	MPLAEVGGFLGGLEGLGQQVGSFLLPPAGERPPLLGERRSAAERSARGGPGAAQLAHLHGILRRRQLYCRTGFHLQILPDGSVQGTRQDHSLEFISVAVGLVSIRGVDSGLYLGMDKGEY GSEKLTSECFREQFEENWYNTYSSNIYKHGDTGRRYFVALNKDGTPRDGARSKRHQKFTHFLPRPVDPERVPELYKDLLMYT with polyhistidine tag at the C-terminus.
Accession	Q9NP95.1
Tag	His-tag at the C-terminus
Mol mass	24.24 kDa
Expiration date	12 months
Bio activity	Measure by its ability to induce 3T3 cells proliferation. The ED50 for this effect is 1.3-3.2 ng/mL. The specific activity of recombinant human FGF-20 is > 2 × 10 ⁵ IU/mg.


Product feature

Purity	> 98% as determined by SDS-PAGE. Ni-NTA chromatography.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
Shipping	Ice bag
Formulation	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 8.0.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

Background

FGF-20 is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF-20 signals through FGFR 2c and 3c, and is expressed during limb and brain development. Recombinant Human FGF-20 is a 23.2 kDa protein containing 209 amino acid residues.

SDS-PAGE



Proce

Procell[®]system

Procell[®]system

Procell[®]system