A Reliable Research Partner in Life Science and Medicine

# Recombinant Rhesus Macaque Angiotensin-Converting Enzyme 2/ACE-2 (C-10His)

Catalog Number: PKSQ050119

Note: Centrifuge before opening to ensure complete recovery of vial contents.

<b>T</b>			
Des	CFI	$\mathbf{n}\mathbf{n}$	nπ
	$\mathbf{c}_{11}$		$\sigma_{\rm LL}$

**Species** Rhesus Macaque

Source HEK293 Cells-derived Rhesus Macaque ACE2/ACE-2 protein Gln18-Val739, with an C-

terminal His

 Mol\_Mass
 85.1 kDa

 Accession
 ACI04564.1

Bio-activity Immobilized Rhesus Macaque ACE-2-His(Cat#PKSQ050119)at 5µg/ml (100 µl/well)

can bind 2019-nCoV S Protein RBD-SD1-mFc(Cat#PKSR030476). The ED50 of Recombinant 2019-nCoV S Protein RBD-SD1-mFc(Cat#PKSR030476) is 16.8 ng/ml.

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \, \text{EU} \, \text{per} \, \mu \text{g} \, \text{of the protein as determined by the LAL method.}$ 

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

**Shipping** This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel


packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 25mM Tris-HCl, 150mM NaCl, 1mM ZnCl<sub>2</sub>,


pH 7.5.

**Reconstitution** Not Applicable

#### Data



> 95 % as determined by reducing SDS-PAGE.



Immobilized Rhesus Macaque ACE-2-His(Cat#PKSQ050119)at 5µg/ml (100 µl/well) can bind 2019-nCoV S Protein RBD-SD1-mFc(Cat#PKSR030476). The ED50 of Recombinant 2019-nCoV S Protein RBD-SD1mFc(Cat#PKSR030476) is 16.8 ng/ml.

# Background

### For Research Use Only

Fax: 1-832-243-6017

# Elabscience®

#### **Elabscience Bionovation Inc.**

A Reliable Research Partner in Life Science and Medicine

Angiotensin-Converting Enzyme 2 (ACE-2) is an integral membrane protein and a zinc metalloprotease of the ACE family, the ACE family includes somatic and germinal ACE. ACE-2 cleaves angiotensins I and II as a carboxypeptidase, ACE-2 converts angiotensin I to angiotensin 1-9, and angiotensin II to angiotensin 1-7. ACE-2 is also able to hydrolyze apelin-13 and dynorphin-13 with high efficiency. ACE-2 can be high expressed in testis, kidney and heart, in colon, small intestine and ovary at moderate levels. Captopril and lisinopril as the classical ACE inhibitor don't inhibit ACE-2 activity. ACE-2 may play an important role in regulating the heart function.

Fax: 1-832-243-6017