

Recombinant Human Interleukin-2/IL-2 Protein (His Tag)

Catalog Number: PKSH032629



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|---------------------|--|
| Species | Human |
| Mol_Mass | 16.4 kDa |
| Accession | P60568 |
| Bio-activity | Measured in a cell proliferation assay using CTLL-2 mouse cytotoxic T cells The specific activity of Recombinant Human IL-2 is $\geq 1 \times 10^7$ IU/mg. |

Properties

| | |
|-----------------------|--|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 0.01 EU per μ g of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 μ m filtered solution of 10mM Sodium Citrate, 4% Mannitol, pH4.0 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-2(IL-2) is an interleukin; a type of cytokine signaling molecule in the immune system; belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells; NK cells; lymphokine-activated killer cells; monocytes; macrophages and oligodendrocytes.

For Research Use Only