EPPIN Polyclonal Antibody

catalog number: E-AB-13218



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Immunogen Synthetic peptide of human EPPIN

Host Rabbit
Isotype IgG


PurificationAffinity purificationConjugationUnconjugated

buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

WB 1:500-1:2000

Data



Western Blot analysis of NIH/3T3 cell, Human testis and Mouse fat tissue using EPPIN Polyclonal Antibody at dilution of 1:400

Calculated-MV:15 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

This gene encodes an epididymal protease inhibitor, which contains both kunitz-type and WAP-type four-disulfide core (WFDC) protease inhibitor consensus sequences. Most WFDC genes are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene is a member of the WFDC gene family and belongs to the telomeric cluste r. The protein can inhibit human sperm motility, and polymorphisms in this gene are associated witHuman, Mouseale infertility. Read-through transcription also exists between this gene and the downstream WFDC6 (WAP four-disulfide core domain 6) gene.

For Research Use Only