


NRF2 Polyclonal Antibody

catalog number: E-AB-68254


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human NRF2
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:200
Data	





Western blot analysis of various lysates using NRF2 Polyclonal Antibody at ARC50393-02 dilution. HeLa cells were treated by MG132 (50 μ M) at 37°C for 90 minutes.

Observed-MW: 97- 100 kDa
Calculated-MW: 65 kDa/66 kDa/67 kDa




Western blot analysis of various lysates using NRF2 Polyclonal antibody at dilution. C6 cells were treated by MG132 (50 μ M) at 37°C for 90 minutes.

Observed-MW: 97- 100 kDa
Calculated-MW: 65 kDa/66 kDa/67 kDa



Western blot analysis of various lysates using NRF2 Polyclonal Antibody at ARC50393-02 dilution. NIH3T3 cells were treated by MG132 (50 μ M) at 37°C for 90 minutes.

Observed-MW: 97- 100 kDa
Calculated-MW: 65 kDa/66 kDa/67 kDa



Immunohistochemistry of paraffin-embedded human breast cancer using NRF2 Polyclonal antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8

Immunohistochemistry of paraffin-embedded human placenta using NRF2 Polyclonal antibody at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

This gene encodes a transcription factor which is a member of a small family of basic leucine zipper (bZIP) proteins. The encoded transcription factor regulates genes which contain antioxidant response elements (ARE) in their promoters; many of these genes encode proteins involved in response to injury and inflammation which includes the production of free radicals. Multiple transcript variants encoding different isoforms have been characterized for this gene.

For Research Use Only