Recombinant Human IA2/PTPRN Protein (aa 607-Asn686 &aa 795-888, His Tag)

Catalog Number: PKSH032555



Note: Centrifuge before opening to ensure complete recovery of vial contents. Description **Species** Human 22.9 kDa Mol Mass Accession O16849 Not validated for activity **Bio-activity Properties** > 90 % as determined by reducing SDS-PAGE. Purity < 1.0 EU per µg of the protein as determined by the LAL method. Endotoxin Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles. Storage This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel Shipping packs. Upon receipt, store it immediately at $< -20^{\circ}$ C. Formulation Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 1mM DTT, 1mM EDTA, pH 8.0. Not Applicable Reconstitution Data

kDa	MK _{aen} ce R
120 90 60	elanse - cience
40	Elaba
30	
20	Habsolt
14	Josoferice
	and the second se

> 90 % as determined by reducing SDS-PAGE.

Background

Receptor-type tyrosine-protein phosphatase-like N (PTPRN) belongs to the protein-tyrosine phosphatase family and receptor class 8 subfamily. PTPRN contains 1 tyrosine-protein phosphatase domain; is expressed in neuroendocrine cells only. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth; differentiation; mitotic cycle; and oncogenic transformation. It implicated in neuroendocrine secretory processes. It may be involved in processes specific for neurosecretory granules; such as their biogenesis; trafficking or regulated exocytosis or may have a general role in neuroendocrine functions. It seems to lack intrinsic enzyme activity; may play a role in the regulation of secretory granules via its interaction with SNTB2. This PTP was found to be an autoantigen that is reactive with insulin-dependent diabetes mellitus (IDDM) patient sera; and thus may be a potential target of autoimmunity in diabetes mellitus.

For Research Use Only