## Recombinant HumanINHB protein (GST, His Tag)

## Catalog Number: PDEH100875

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	E.coli-derived Human INHB protein Gly311-Ser426, with an N-terminal GST & C-
	terminal His
Calculated MW	37.7 kDa
Observed MW	36 kDa
Accession	P08476
Bio-activity	Not validated for activity
Properties	
Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}C$ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with 5% Trehalose and 5%
	Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of
	0.5 mg/mL. Concentration is measured by UV-Vis.

## Background

Activin and inhibin are two closely related protein complexes that have almost directly opposite biological effects. Activins, members of the TGF-beta superfamily, are disulfide-linked dimeric proteins originally purified from gonadal fluids as proteins that stimulated pituitary follicle stimulating hormone (FSH) release. Inhibins/activins are involved in regulating a number of diverse functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition. Activins are homodimers or heterodimers of the various beta subunit isoforms, while inhibins are heterodimers of a unique alpha subunit and one of the various beta subunits.