

Recombinant Human SIGIRR/TIR8 Protein (Fc Tag)

Catalog Number: PKSH030884

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|---|
| Species | Human |
| Source | HEK293 Cells-derived Human SIGIRR/TIR8 protein Met 1-His118, with an C-terminal hFc |
| Calculated MW | 39.5 kDa |
| Observed MW | 47-54&33 kDa |
| Accession | Q6IA17-1 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|--|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Single Ig IL-1-related receptor (SIGIRR) or TIR8 is a member of Toll-like receptor-interleukin 1 receptor signaling (TLR-IL-1R) receptor superfamily. Although SIGIRR/TIR8 shows the typical conserved motifs that characterize the IL-1R and Toll superfamily; it is structurally and functionally distinct from both. SIGIRR/TIR8 has only one Ig domain in its extracellular portion whereas the IL-1R family contains three Ig folds. An unusually long cytoplasmic domain is reminiscent of the structure of drosophila Toll; yet the SIGIRR peptide sequence is more closely related to IL-1RI. SIGIRR/TIR8 was mainly expressed in mouse and human epithelial tissues such as kidney; lung and gut. Resting and activated T and B lymphocytes and monocytes-macrophages expressed little or no SIGIRR/TIR8; with the exception of the mouse GG2EE macrophage line. Inflammation is enhanced in SIGIRR-deficient mice. SIGIRR negatively modulates immune responses. Inflammation is enhanced in SIGIRR-deficient mice; as shown by their enhanced chemokine induction after IL-1 injection and reduced threshold for lethal endotoxin challenge.