Recombinant Human CLIC2 Protein (His Tag)

Catalog Number: PKSH032247



Note: Centrifuge before opening to ensure complete recovery of vial contents.

				ption		
	06		TO T	m	11	\mathbf{n}
v					ш	w

 Species
 Human

 Mol_Mass
 30.5 kDa

 Accession
 O15247

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu\text{g of the protein as determined by the LAL method.}$

Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel


packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 100mM NaCl, 1mM DTT,

20% Glycerol, pH 8.0.

Reconstitution Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Chloride Intracellular Channel Protein 2 (CLIC2) is a critical component of all living cells; it regulatescellular traffic of Chloride ion and it can be inserted into membranes and form chloride ion channels. Membrane insertion seems to be redox-regulated and may occur only under oxydizing conditions, channel activity depends on the pH. CLIC2 is involved in regulating membrane potential and organic solute transport. CLIC2 modulates the activity of RYR2 and inhibits Calcium influx. CLIC2 can be detected in the adult brain, liver, lung, heart, stomach, spleen and testis. It is expressed in fetal liver and adult skeletal muscle. CLIC2 is a potential candidate for one of many diseases linked to Xq28.

For Research Use Only