

Recombinant Mouse APRIL/TNFSF13 Protein(Fc Tag)

Catalog Number: PDMM100178

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	Mammalian-derived Mouse APRIL/TNFSF13 proteins Ala96-Leu241, with an C-terminal Fc
Calculated MW	40.9 kDa
Observed MW	42 kDa
Accession	Q9D777
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse APRIL/TNFSF13 proteins, 2 µg/lane of Recombinant Mouse APRIL/TNFSF13 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 42 KD

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7

TNFSF13 is a member of the tumor necrosis factor (TNF) ligand family. It is a ligand for TNFRSF17/BCMA. TNFSF13 is lowly expressed in normal tissues, but is elevated in several types of tumors and transformed cell lines. It is important for B cell development. TNFSF13 may also play a role in T-independent type II antigen responses and T cell survival, and induce proliferation/survival of non lymphoid cells. It exists as a functional homotrimer. It can bind to two cell surface receptors, BCMA and TACI, which it shares with BAFF to exert downstream T-and B-cell regulatory effects. TNFSF13 also has been demonstrated to bind to proteoglycans on the cell surface.