



A Reliable Research Partner in Life Science and Medicine

PE/Cyanine7 Anti-Human CD18 Antibody[TS1/18.1.2.11]

Catalog Number: E-AB-F1057H

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| _ | | 4.0 | |
|---|-------|------|-----|
| | escri | MTI. | on. |
| ш | COUL | | UII |

Reactivity Human Host Mouse

 Isotype
 Mouse IgG1, κ

 Clone No.
 TS1/18.1.2.11

Isotype Control PE/Cyanine7 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792H]

Conjugation PE/Cyanine 7

Conjugation Information PE/Cyanine7 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 775 nm

(e.g., a 780/60 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD18;Cell surface adhesion glycoproteins LFA-1/CR3/p150+95 subunit beta;

Complement receptor C3 subunit beta;Integrin beta-2;Itgb2

 Uniprot ID
 P05107

 Gene ID
 3689

Background CD18 is a 90-95 kD type I transmembrane protein also known as integrin β2 subunit,

LFA-1 β subunit, and $\beta2$ integrin. CD18 non-covalently associates with CD11a, CD11b

or CD11c. CD18 is expressed on all leukocytes. CD18 and associated $\boldsymbol{\alpha}$ chains

function in adhesion and signaling in hematopoietic cells.

Web: www.elabscience.cn