

## Recombinant Human IL17RA Protein (His Tag)

**Catalog Number:** PKSH031368

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human IL17RA protein Met 1-Trp 320, with an C-terminal His
<b>Calculated MW</b>	35.0 kDa
<b>Accession</b>	NP_055154.3
<b>Bio-activity</b>	Measured by its ability to inhibit IL-17-induced IL-6 secretion by NIH- 3T3 mouse embryonic fibroblast cells. The ED <sub>50</sub> for this effect is 0.2- 1.2µg/mL in the presence of 5 ng/mL recombinant human IL- 17.


### Properties

<b>Purity</b>	> 97 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 97 % as determined by reducing SDS-PAGE.



Measured by its ability to inhibit IL-17-induced IL-6 secretion by NIH- 3T3 mouse embryonic fibroblast cells. The ED<sub>50</sub> for this effect is 0.2- 1.2µg/mL in the presence of 5 ng/mL recombinant human IL- 17.

### Background

#### For Research Use Only

Interleukin-17 receptor (IL-17R); also known as Interleukin-17 receptor A (IL-17RA) and CD217 antigen (CD217); is a cytokine receptor which binds interleukin 17. IL-17R/IL-17RA (CD217) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. IL-17R/IL-17RA (CD217) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor IL-17RA play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors; this receptor likely has a multimeric structure. Defects in IL-17R/IL-17RA (CD217) are the cause of familial candidiasis type 5 (CANDF5). CANDF5 is a rare disorder with altered immune responses and impaired clearance of fungal infections; selective against *Candida*. It is characterized by persistent and/or recurrent infections of the skin; nails and mucous membranes caused by organisms of the genus *Candida*; mainly *Candida albicans*.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017