

A Reliable Research Partner in Life Science and Medicine

PHF21A Polyclonal Antibody

catalog number: E-AB-18538

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Immunogen Fusion protein of human PHF21A

Host Rabbit Isotype IgG


Purification Antigen affinity purification

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


Applications Recommended Dilution

WB 1:500-1:2000 **IHC** 1:40-1:200

Data




Western blot analysis of 293T Hela and A375 cell lysates using PHF21A Polyclonal Antibody at dilution of 1:400



Immunohistochemistry of paraffin-embedded Human prost at e cancer tissue using PHF21A Polyclonal Antibody at dilution of 1:50(×200)

Observed-MW:Refer to figures

Calculated-MW:75 kDa



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using PHF21A Polyclonal Antibody at dilution of 1:50(×200)

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack,upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. PHF21A (PHD finger protein 21A), also known as BRAF35-HDAC complex protein BHC80, is a 680 amino acid nuclear protein that contains one PHD-type zinc finger and one A.T hook DNA-binding domain, suggesting involvement in transcriptional regulation events. PHF21A is a component of the BHC complex, which is responsible for repressing transcription of neuron-specific genes in non-neuronal cells. The BHC complex acts as a chromatin modifier that deacetylates and demethylates specific sites on histones. PHF21A may act as a scaffold within the BHC complex. Predominantly expressed in brain, three isoforms of PHF21A exist as a result of alternative splicing events.

For Research Use Only

Toll-free: 1-888-852-8623 Web:<u>w w w .elabscience.com</u>

Tel: 1-832-243-6086 Email:techsupport@elabscience.com Fax: 1-832-243-6017