

A Reliable Research Partner in Life Science and Medicine

## **PHF21A Polyclonal Antibody**

catalog number: E-AB-18538

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Mouse

**Immunogen** Fusion protein of human PHF21A

Host Rabbit Isotype IgG


**Purification** Antigen affinity purification

**Buffer** Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

## **Applications** Recommended Dilution

**WB** 1:500-1:2000 **IHC** 1:40-1:200

#### Data




Western blot analysis of 293T Hela and A375 cell lysates using PHF21A Polyclonal Antibody at dilution of 1:400

# Immunohistochemistry of paraffin-embedded Human prost at e cancer tissue using PHF21A Polyclonal Antibody at dilution of 1:50(×200)



#### Calculated-MV:75 kDa



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using PHF21A Polyclonal Antibody at dilution of 1:50(×200)

## Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

## Background

## For Research Use Only

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. PHF21A (PHD finger protein 21A), also known as BRAF35-HDAC complex protein BHC80, is a 680 amino acid nuclear protein that contains one PHD-type zinc finger and one A.T hook DNA-binding domain, suggesting involvement in transcriptional regulation events. PHF21A is a component of the BHC complex, which is responsible for repressing transcription of neuron-specific genes in non-neuronal cells. The BHC complex acts as a chromatin modifier that deacetylates and demethylates specific sites on histones. PHF21A may act as a scaffold within the BHC complex. Predominantly expressed in brain, three isoforms of PHF21A exist as a result of alternative splicing events.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web:www.elabscience.com
 Email:techsupport@elabscience.com