Recombinant Human CD200R1 Protein (Fc Tag)

Catalog Number: PKSH032207



Human
53.9 kDa
Q8TD46
Not validated for activity
> 95 % as determined by reducing SDS-PAGE.
< 1.0 EU per µg of the protein as determined by the LAL method.
Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
This product is provided as lyophilized powder which is shipped with ice packs.
Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
before lyophilization.
Please refer to the specific buffer information in the printed manual.
Please refer to the printed manual for detailed information.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

kDa	MK	R
120 90 60	=	-
40	-	New Der
30	-	Part State
20	-	
14	-	1
		the strength of the strength o

> 95 % as determined by reducing SDS-PAGE.

Background

Cell surface glycoprotein CD200 Receptor 1 (CD200R1) is the receptor for the CD200 (OX-2) membrane glycoprotein. CD200R1 contains one C2- type Ig-like domain and one V-type Ig-like domain within its extracellular domain and a PTBsignaling motif in cytoplasmic domain. CD200R1 and CD200 associate via their respective N-terminal Ig-like domains. CD200R1 is restricted primarily to mast cells, basophils, macrophages, and dendritic cells. It propagates inhibitory signals despite its lacking a cytoplasmic ITIM (immunoreceptor tyrosinebased inhibitory motif). The receptor-substrate interaction may function as a myeloid downregulatory signal.

For Research Use Only