



A Reliable Research Partner in Life Science and Medicine

Canine CRP Antibody Pair Set

Catalog No. E-KAB-0396 Applications ELISA

Synonyms PTX1;Pentraxin 1

Kit components & Storage

Title	Specifications	Storage
Canine CRP Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year. Avoid
		freeze/thaw cycles.
Canine CRP Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C for one year. Avoid
		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0396)	
		Canine CRP Capture Antibody	Canine CRP Detection Antibody
			(Biotin)
Immunogen	Immunogen	Natural Canine CRP protien	Natural Canine CRP protien
Information	Swissprot	T2KEN6	
Product details	Reactivity	Canine	Canine
	Host	Mouse	Mouse
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Protein A or G	Protein A or G
	Specificity	Detects Canine CRP in ELISAs.	·

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Canine CRP Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Canine CRP Capture	
Capture		Antibody	10
			Alse
EL IGA	1 1000 1 1000	G : GDD D : i	Optical Density
ELISA	1:1000-1:10000	Canine CRP Detection	8 .
Detection		Antibody (Biotin)	0.1
			1 10 100 1000
			Canine CRP Concentration (ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The protein encoded by this gene belongs to the pentaxin family. It is involved in several host defense related functions based on its ability to recognize foreign pathogens and damaged cells of the host and to initiate their elimination by interacting with humoral and cellular effector systems in the blood. Consequently, the level of this protein in plasma increases greatly during acute phase response to tissue injury, infection, or other inflammatory stimuli.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com