

Recombinant Human CXCL2/MIP-2 Protein

Catalog Number:PKSH033705



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	C-X-C Motif Chemokine 2;Growth-Regulated Protein Beta;Gro-Beta;Macrophage Inflammatory Protein 2-Alpha;MIP2-Alpha;CXCL2;GRO2;GROB;MIP2A;SCYB2;GRO beta;MIP2
Species	Human
Expression Host	E.coli
Sequence	Val 35-Asn107
Accession	P19875
Calculated Molecular Weight	8.7 kDa
Observed molecular weight	12 kDa
Tag	N-His
Bioactivity	Measure by its ability to chemoattract human PBMCs using a concentration range of 10.0 - 100.0 ng/mL. Note: Results may vary from different PBMC donors.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Chemokine Ligand 2 (CXCL2) is a small secreted cytokine which belongs to the CXC chemokine family. It is secreted by monocytes and macrophages and chemotactic for polymorphonuclear leukocytes and hematopoietic stem cells. CXCL2 mobilizes cells by interacting with a cell surface chemokine receptor called CXCR2. It has been known to regulate immune functions mainly by chemo-attracting neutrophils. It is produced by activated monocytes and neutrophils and

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CXCL2/MIP-2 Protein

Catalog Number:PKSH033705



expressed at sites of inflammation. It is a hematoregulatory chemokine, which suppresses hematopoietic progenitor cell proliferation. It can be induced by receptor activator of NF-kappaB ligand, the osteoclast (OC) differentiation factor, through JNK and NF-kappaB signaling pathways in OC precursor cells. CXCL2 in turn enhanced the proliferation of OC precursor cells of bone marrow-derived macrophages (BMMs) through the activation of ERK. Knockdown of CXCL2 inhibited both the proliferation of and the ERK activation in BMMs. During osteoclastogenesis CXCL2 stimulated the adhesion and the migration of BMMs. CXCL2 is a novel therapeutic target for inflammatory bone destructive diseases.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017