

## CCL15/MIP-1δ Polyclonal Antibody(Capture/Detector)

catalog number: AN004500P

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human CCL15/MIP-1δ Protein expressed by E.coli
<b>Host</b>	Rabbit
<b>Isotype</b>	Rabbit IgG
<b>Purification</b>	Antigen Affinity Purification
<b>Conjugation</b>	Unconjugated
<b>Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

### Applications

### Recommended Dilution

<b>ELISA Capture</b>	2-8 µg/mL
<b>ELISA Detector</b>	0.1-0.4 µg/mL

### Data



Sandwich ELISA-Recombinant Human CCL15/MIP-1δ Protein standard curve. Background subtracted standard curve using Anti-CCL15/MIP-1δ antibody(AN004500P) (Capture), Anti-CCL15/MIP-1δ antibody(AN004500P) (Detector). The reference range value is 15.6 - 1,000 pg/mL for human.

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

### Background

Chemokine ligand 15 (CCL15), also known as leukotactin-1, MIP5, MIP1 and HCC-2, is a small cytokine belonging to the C-C chemokine family. CCL15 is prevalently expressed in liver, small intestine, colon, and in certain leukocytes and macrophages of the lung. It is chemotactic for neutrophils, monocytes, and lymphocytes and elicits its effects by binding to cell surface chemokine receptors like CCR1 and CCR3.

### For Research Use Only