Recombinant Latexin/LXN Monoclonal Antibody

catalog number: AN300180P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human
Immunogen	Recombinant Human Latexin / LXN Protein
Host	Rabbit
Isotype	IgG
Clone	9B8
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS
Applications	Recommended Dilution
WB	1:500-1:2000
FCM	1:25-1:100
ICC/IF	1:20-1:100
IP	4-6 μL/mg of lysate

Data



Flow cytometric analysis of Human LXN expression on

MCF-7 cells. The cells were stained with purified anti-Human LXN, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.





Immunofluorescence analysis of Human LXN in MCF7 cells. Cells were fixed with 4% PFA, permeabilzed with 1% Triton X-100 in PBS, blocked with 10% serum, and incubated with Rabbit anti-Human LXN Monoclonal Antibody (1:60) at 4°C overnight. Then cells were stained with the Alexa Fluor® 488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI for nuclear staining (blue). Positive staining was localized to



For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com Tel: 1-832-243-6086 Email:techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc. A Reliable Research Partner in Life Science and Medicine

Immunoprecipitation analysis using 2 μL anti-LXN Monoclonal Antibody and 15 μl of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using LXN Monoclonal Antibody at a dilution of 1:200. Lane A:0.5 mg A549 Whole Cell Lysate, Lane B:0.5 mg Hela Whole Cell Lysate **Observed-MW:25 kDa**

Western Blot with Latexin / LXN Monoclonal Antibody at dilution of 1:500. Lane A: A549 Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:25 kDa Calculated-MW:25 kDa

Calculated-	MW:25 kDa
Preparation & Storage	
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag
Background	

Latexin, also known as endogenous carboxypeptidase inhibitor, tissue carboxypeptidase inhibitor, TCI, ECI, and LXN, is a cytoplasm protein that belongs to the protease inhibitor I47 (latexin) family. It is highly expressed in the heart, prostat e, ovary, kidney, pancreas, and colon. Latexin/LXN is the only known endogenous specific inhibitor of zinc-dependent metallocarboxypeptidases (MCPs) present in mammalians so far. Latexin is originally identified as a molecular marker for the regional specification of the neocortex in development in rats. The 222 amino acid latexin in the human shows different expression distribution with high levels in heart, prostate, ovary, kidney, pancreas, and colon, but only moderate or low levels in other tissues including the brain. Latexin is also expressed at high levels and is inducible in macrophages in concert with other protease inhibitors and potential protease targets, and thus is suggested to play a role in inflammation and innate immunity pathways. Despite the non-detectable sequence similarity with plant and parasite inhibitors, Latexin is related to a human putative tumor suppressor protein, TIG1. Also, Latexin is implicated in Alzheimer's disease.