

Human PDGF-BB Antibody Pair Set

Catalog No. E-KAB-0495

Applications

ELISA

Synonyms PDGFBB

Kit components & Storage

Title	Specifications	Storage
Human PDGF-BB Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze/thaw cycles.
Human PDGF-BB Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0495)	
		Human PDGF-BB Capture Antibody	Human PDGF-BB Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human PDGF-BB protien	Recombinant Human PDGF-BB protien
	Swissprot	P01127	
Product details	Reactivity	Human	Human
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50% glycerol; pH 7.5	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.5
	Purify	Antigen Affinity	Antigen Affinity
Specificity	Detects Human PDGF-BB in ELISAs.		

For Research Use Only

Applications

Human PDGF-BB Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4 µg/mL	Human PDGF-BB Capture Antibody	
ELISA Detection	1:1000-1:10000	Human PDGF-BB Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The protein encoded by this gene is a member of the platelet-derived growth factor family. The four members of this family are mitogenic factors for cells of mesenchymal origin and are characterized by a motif of eight cysteines. This gene product can exist either as a homodimer (PDGF-BB) or as a heterodimer with the platelet-derived growth factor alpha polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds. Mutations in this gene are associated with meningioma. Reciprocal translocations between chromosomes 22 and 7, at sites where this gene and that for COL1A1 are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans resulting from unregulated expression of growth factor. Two alternatively spliced transcript variants encoding different isoforms have been identified for this gene.