

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 647 Anti-Mouse CD279/PD-1 Antibody[29F.1A12]

Catalog Number: E-AB-F1131M

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Rat Host

Isotype Rat IgG2a, ĸ Clone No. 29F.1A12

Isotype Control Elab Fluor® 647 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832M]

Conjugation Elab Fluor®647

Conjugation Information Elab Fluor® 647 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).


Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Con-A stimulated C57BL/6 splenocytes (3 days) are stained

with Elab Fluor® 647 Anti-Mouse CD279/PD-1 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Keep as concentrated solution. **Storage**

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names Programmed Death-1;PD-1

Uniprot ID Q02242 Gene ID 18566

For Research Use Only

Fax: 1-832-243-6017 Tel: 1-832-243-6086 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

CD279, also known as programmed death-1 (PD-1), is a 50-55 kD glycoprotein belonging to the CD28 family of the Ig superfamily. PD-1 is expressed on activated splenic T and B cells and thymocytes. It is induced on activated myeloid cells as well. PD-1 is involved in lymphocyte clonal selection and peripheral tolerance through binding its ligands, B7-H1 (PD-L1) and B7-DC (PD-L2). It has been reported that PD-1 and PD-L1 interactions are critical to positive selection and play a role in shaping the T cell repertoire. PD-L1 negative costimulation is essential for prolonged survival of intratesticular islet allografts.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Email:techsupport@elabscience.com Web:www.elabscience.com